

DOSAGE & ADMINISTRATION

Dosage of MSUD express to be determined by the clinician or dietitian.

1. The daily protein requirement is given as MSUD express (100g is equivalent to 60g of protein).
2. Additional protein is provided from natural protein sources in medically prescribed quantities to provide Leucine, Isoleucine and Valine in controlled amounts which is essential for normal development.
3. Foods high in carbohydrate and fat but low in protein should be provided (as an energy source) to supply the remainder of the calories.
4. A supplement of Isoleucine and Valine is given if required (if plasma concentrations fall too low).

PREPARATION GUIDELINES

Place the contents of the sachet into the beaker provided, add a FlavourPac sachet if required and fill with cold water to about half way up the beaker approximately 80ml of water. Secure the lid and shake for about 10 seconds. Drink immediately.

Water or permitted drinks should be taken after the MSUD express.

The product should be freshly prepared and when possible used immediately. However, any unused portions should be refrigerated and used within 24 hours - re-shake before use.

IMPORTANT NOTICE

Must only be given to patients with proven Maple Syrup Urine Disease and under strict medical supervision.

Not for intravenous use.

Not recommended as a sole source of nutrition.

Contains a source of phenylalanine.

Not recommended for children under 8 years of age.

STORAGE

Store in a cool, dry place.

PACK SIZE

MSUD express is available in packs of 30 x 25g sachets.

ACBS and GMS APPROVED



Innovation in Nutrition

AUSTRALIA · GERMANY · IRELAND · ITALY
SCANDINAVIA · UNITED KINGDOM · USA

Manufactured in the UK for VitaFlo International Ltd.
11-16 Century Building, Brunswick Business Park,
Liverpool L3 4BL

Tel: 0151 709 9020 Fax: 0151 709 9727

E-mail: vitaflo@vitaflo.co.uk Web: www.vitaflo.co.uk

VitaFlo and MSUD express are Trade Marks of
VitaFlo International Ltd.

Code: 0404